

Large B -Fields and Noncommutative Solitons¹

Jorge Moreno²

*Departamento de Física
Centro de Investigación y de Estudios Avanzados del I.P.N.
Apdo. Postal 14-740, 07000, México D.F. México*

Abstract. The purpose of this talk is to review a few issues concerning noncommutativity arising from String Theory. In particular, it is shown how in Type IIB Theory, the annihilation of a $D3 - \overline{D3}$ brane pair yields a $D1$ -string. This object, in the presence of a large B -field and fermions, happens to be a complex noncommutative soliton endowed with superconductivity.

INTRODUCTION

Type IIB Superstring Theory allows two interesting ingredients: stable-BPS RR-charged Dp -branes (p odd) and a massless antisymmetric $B_{\mu\nu}$ field. Although a $D3 - D3$ brane system is stable; a $D3 - \overline{D3}$ case is not, due to the presence of a tachyon in its spectrum [1]. The job of the B -field in the low energy limit is to introduce noncommutativity [2]. In the $D3 - \overline{D3}$ brane configuration, we may turn on a large B -field along two worldvolume spatial coordinates. The effect of this is that the complex tachyon allows a GMS soliton [3]. This object appears to be superconducting in the presence of fermions arising from the open string sector [4].

B -FIELDS AND NONCOMMUTATIVITY

Consider an open string attached to a D -brane. The OPE has the form

$$e^{ik_1 \cdot X} e^{ik_2 \cdot X} \sim (\tau - \tau')^{2\alpha' g^{\mu\nu} k_{1\mu} \cdot k_{2\nu}} \times e^{i(k_1+k_2) \cdot X} + \dots \quad (1)$$

However, the introduction of a large B -field alters this to

$$e^{ik_1 \cdot X} e^{ik_2 \cdot X} \sim (\tau - \tau')^{2\alpha' G^{\mu\nu} k_{1\mu} \cdot k_{2\nu}} \times \left[e^{-\frac{i}{2} \Theta^{\mu\nu} k_{1\mu} \cdot k_{2\nu}} \right] \times e^{i(k_1+k_2) \cdot X} + \dots, \quad (2)$$

¹⁾ This work is dedicated to my parents Juan Moreno and Martha Soto.

²⁾ E-mail address: jmoreno@fis.cinvestav.mx

where $G^{\mu\nu} = \left(\frac{1}{g+2\pi\alpha'B}g\frac{1}{g-2\pi\alpha'B}\right)^{\mu\nu}$ is the effective metric seen by the open string modes, and $\Theta^{\mu\nu} = -(2\pi\alpha')^2\left(\frac{1}{g+2\pi\alpha'B}B\frac{1}{g-2\pi\alpha'B}\right)^{\mu\nu}$ is the noncommutativity parameter matrix [3]. Likewise, the new term in brackets is known -in configuration space- as the Moyal $*$ product. Thus, in the low energy effective theory, fields get $*$ -multiplied.

NONCOMMUTATIVE SOLITONS

The idea of GMS solitons was cleverly used in [5] to construct real solutions to the tachyon in bosonic D -branes. In this work, complex tachyons in RR D -brane pairs are considered instead. For a $D3 - \overline{D3}$ brane pair in the presence of a large B -field along the $x - y$ plane, the action of the tachyon is

$$S^{(\Sigma_{3+1})} = \int_{\Sigma_{3+1}} dx dy dz dt \left[\overline{D^\mu T} * D_\mu T - V_* (T, \overline{T}) \right], \quad (3)$$

where $[x, y] = i\theta$, and z and t are commutative coordinates. Also, $D_\mu T = \partial_\mu T - iA_\mu * T + i\widetilde{A}_\mu * T$, where A_μ and $\widetilde{A}_\mu$ are the respective gauge fields in each of the D -brane's Chan-Paton $U(1)$ symmetry groups.

We make three assumptions:

- The potential is a polynomial:

$$V_* (T, \overline{T}) = \sum_{k=1}^n a_k (\overline{T} * T)^k. \quad (4)$$

- The gauge field $R_\mu \equiv A_\mu - \widetilde{A}_\mu$ has the form:

$$R_\mu = R_\mu(z, t). \quad (5)$$

- We'll focus on solutions of the form:

$$T = T(x, y), \quad \overline{T} = \overline{T}(x, y). \quad (6)$$

Eventually, it is shown that the solution is

$$T = t_* T_0, \quad \overline{T} = \overline{t}_* \overline{T}_0, \quad (7)$$

where t_* and $\overline{t}_*$ solve the equations of motion in the commutative case, while $T_0, \overline{T}_0$ satisfy

$$(\overline{T}_0 * T_0)^k = \overline{T}_0 * T_0. \quad (8)$$

Since $[x, y] = i\theta$ is analogous to $[\hat{q}, \hat{p}] = i$ in quantum mechanics, we identify $T_0 \leftrightarrow i|0\rangle\langle 0|$ and $\overline{T}_0 \leftrightarrow -i|0\rangle\langle 0|$ in the Simple Harmonic Oscillator basis.

Applying the Weyl-Wigner-Moyal correspondence yields the following lowest energy solitonic solution:

$$T(x, y) = 2it_*e^{-r^2}, \quad \overline{T}(x, y) = -2i\overline{t}_*e^{-r^2}, \quad (9)$$

where $r^2 = x^2 + y^2$ [4].

THE NONCOMMUTATIVE SUPERCONDUCTING STRING

Given a $(3+1)$ Dirac spinor Ψ , with two-component entries ψ_R and ψ_L obeying $\vec{\sigma} \cdot \hat{p}\psi_R = \psi_R$ and $\vec{\sigma} \cdot \hat{p}\psi_L = -\psi_L$, the action for the spinor coupled to the complex soliton has the form:

$$S^{(\Sigma_{3+1})} = \int_{\Sigma_{3+1}} dt dz dx dy \left[f(\overline{T}) * \overline{\Psi} * g(T) \not{D}\Psi \right], \quad (10)$$

where f and g are polynomials, and $\not{D}\Psi = \gamma^\mu (\partial_\mu \Psi - iR_\mu * \Psi)$.

In terms of operators, we may express our spinors as [6]:

$$\hat{\psi}_{L,R}(x^\mu) = \sum_{m,n \geq 0} \psi_{mn}^{L,R}(z, t) |m\rangle \langle n|. \quad (11)$$

In order to find the effective theory along the *string* (the z, t coordinates,) we make two assumptions:

- $\theta \longrightarrow \infty$, which means that the noncommutative kinetic part is negligible.
- $\psi \longrightarrow \psi^L$ (as in Witten's superconducting string [7]).

Therefore, the action for the *Noncommutative D-string* is

$$S^{(\Sigma_{1+1})} = -2\pi i \theta f(\overline{t}_*) g(t_*) \int_{\Sigma_{1+1}} \left[i\overline{\psi}^L \sigma^a D_a \psi^L - \overline{\psi}^L m \psi^L \right], \quad (12)$$

where ψ^L denotes ψ_{00}^L in (11), m is a “mass” matrix ($m(z, t) = \sigma^\alpha R_\alpha$)³.

SUPERCONDUCTIVITY

In the massless case (after rescaling the action and getting rid of the unnecessary L subscript)

³⁾ $a = z, t$ and $\alpha = x, y$.

$$S^{(\Sigma_{3+1})} = \int_{\Sigma_{3+1}} dz dt \left(i \bar{\psi} \sigma^a D_a \psi \right). \quad (13)$$

According to the bosonization technique [7], in two dimensions we can equivalently describe the theory by either bosons or fermions.

This is done by introducing a scalar field $\zeta(z, t)$ such that

$$\bar{\psi} \sigma^a \psi = \frac{1}{\sqrt{\pi}} \varepsilon^{ab} \partial_b \zeta. \quad (14)$$

It can be shown that the kinetic term corresponds to

$$i \bar{\psi} \sigma^a D_a \psi = \frac{1}{2} (\partial_a \zeta) (\partial^a \zeta) - \frac{1}{\sqrt{\pi}} R_a \varepsilon^{ab} \partial_b \zeta, \quad (15)$$

which is associated to a conserved current

$$J^a = \partial^a \zeta + \frac{1}{\sqrt{\pi}} \varepsilon^{ab} R_b. \quad (16)$$

However, this current may be expressed in terms of another scalar:

$$J^a = \varepsilon^{ab} \partial_b \varphi. \quad (17)$$

Thus, $\partial_b \varphi = -\varepsilon_{ba} J^a$ and

$$\partial^b \partial_b \varphi = -\partial^b \varepsilon_{ba} \left(\partial^a \zeta + \frac{1}{\sqrt{\pi}} \varepsilon^{ab} R_b \right) = -\frac{1}{\sqrt{\pi}} \partial^b R_b. \quad (18)$$

This means that from $\partial_a \varphi = -\varepsilon_{ac} J^c$, we get that $-\varepsilon_{ac} \partial^a J^c = -\frac{1}{\sqrt{\pi}} \partial^a R_a$. In other words, for J^3 (the current along the string):

$$\frac{dJ^3(z, t)}{dt} = \frac{1}{\sqrt{\pi}} \frac{dR_0(z, t)}{dz}, \quad (19)$$

which has the following solution:

$$J^3(z, t) = \frac{1}{\sqrt{\pi}} [R_0(z, \tau) - R_0(z, \tau_i)]. \quad (20)$$

This means that the current is nondecaying as long as the gauge fields has different values when it's turn on ($t = \tau_i$) and turn off ($t = \tau$).

ACKNOWLEDGMENTS

It is a pleasure to thank my advisor Dr. H. García-Compeán for useful discussions and enjoyable collaboration. I'm very grateful to J.M.L. Bauche A. for helping me write this manuscript. I also thank the organizers of the *VIII Mexican Workshop on Particles and Fields* for their hospitality. This work was supported in part by the CONACyT Fellowship No. 33951E.

REFERENCES

1. J. H. Schwarz, “TASI Lectures on Non-BPS D-Brane Systems”, hep-th/9908144.
2. R. Gopakumar, S. Minwalla, and A. Strominger, “Noncommutative Solitons,” JHEP **0005** (2000) 020, hep-th/0003160.
3. N. Seiberg and E. Witten, “String Theory and Noncommutative Geometry,” JHEP **09** (1999) 032, hep-th/9908142.
4. H. García-Compeán and J. Moreno, “Remarks on Noncommutative Solitons,” hep-th/0110119.
5. J. Harvey, P. Kraus, F. Larsen, and E.J. Martinec, “D-branes and Strings as Noncommutative Solitons,” JHEP **0007** (2000) 042, hep-th/0005031.
6. L. Pilo and A. Riotto, “The Non-commutative Brane World,” JHEP **03** (2001) 015, hep-ph/0012174
7. E. Witten, “Superconducting Strings,” Nucl. Phys. B **249** (1985) 557.